



# GS-323

VI Semester B.A./B.Sc. Examination, May/June - 2019

## MATHEMATICS

### Mathematics - VII

(CBCS) (F+R) (2016-17 & Onwards)

Time : 3 Hours

Max. Marks : 70

**Instructions :** Answer *all* questions.

### PART - A

Answer **any five** sub-questions.

**5x2=10**

1. (a) In a vectorspace  $V(F)$  show that  $C(-\alpha) = -(C\alpha)$ ,  $\forall C \in F, \alpha \in V$
- (b) Prove that the set  $S = \{(3, 2, -1), (0, 4, 5), (6, 4, -2)\}$  is Linearly dependent in  $V_3(\mathbb{R})$ .
- (c) Find the matrix of the linear transformation  
 $T : V_2(\mathbb{R}) \rightarrow V_2(\mathbb{R})$  defined by  
 $T(x, y) = (2x+3y, 4x-5y)$  with respect to standard bases.
- (d) Define Rank and Nullity of linear transformation.
- (e) In a cylindrical coordinate system prove that  $\hat{e}_\phi \cdot \hat{e}_z = 0$
- (f) Solve  $\frac{x dx}{y^2 z} = \frac{dy}{zx} = \frac{dz}{y^2}$
- (g) Form the partial differential equation by eliminating the arbitrary constants from  $2Z = \frac{x^2}{a^2} + \frac{y^2}{b^2}$
- (h) Solve  $\sqrt{p} + \sqrt{q} = 1$

**P.T.O.**



## PART - B

Answer **any two full** questions.

**2x10=20**

2. (a) A Subset  $W$  of a vectorspace  $V(F)$  is a subspace if and only if
- $\alpha, \beta \in W \Rightarrow \alpha + \beta \in W$
  - $C \in F, \alpha \in W \Rightarrow C \cdot \alpha \in W$
- (b) Find the basis and dimension of the subspace spanned by  $(2, 4, 2)$ ,  $(1, -1, 0)$ ,  $(1, 2, 1)$ ,  $(0, 3, 1)$  in  $V_3(\mathbb{R})$

**OR**

3. (a) A set of non zero vectors  $(\alpha_1, \alpha_2, \dots, \alpha_n)$  of vectorspace  $V(F)$  is linearly dependent if and only if one of these vectors say  $\alpha_k$  ( $2 \leq k \leq n$ ) is expressed as a linear combination of its preceding ones.
- (b) Show that the subset  $W = \{(x_1, x_2, x_3) / x_1 + x_2 + x_3 = 0\}$  is a subspace of  $V_3(\mathbb{R})$
4. (a) If  $T : U \rightarrow V$  is a linear transformation then prove that.
- $T(0) = 0'$ , where  $0$  and  $0'$  are the zero vectors of  $U$  and  $V$  respectively.
  - $T(-\alpha) = -T(\alpha), \forall \alpha \in U$
- (b) Verify whether  $T : V_2(\mathbb{R}) \rightarrow V_2(\mathbb{R})$  is a linear transformation defined by  $T(x, y) = (3x + 2y, 3x - 4y)$

**OR**

5. (a) Find the range space, null space, rank, nullity and hence verify rank nullity theorem for  $T : V_3(\mathbb{R}) \rightarrow V_3(\mathbb{R})$  given by  $T(x, y, z) = (x + y, x - y, 2x + z)$
- (b) Show that the linear transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $T(e_1) = e_1 + e_2$ ,  $T(e_2) = e_1 + e_3$ ,  $T(e_3) = e_1 + e_2 + e_3$  is non-singular where  $\{e_1, e_2, e_3\}$  is the standard basis of  $\mathbb{R}^3$ .

## PART - C

Answer **any two full** questions.

**2x10=20**

6. (a) Verify the condition for integrability and solve  $(2x^2 + 2xy + 2xz^2 + 1) dx + dy + 2z dz = 0$
- (b) Solve  $p \tan x + q \tan y = \tan z$

**OR**

7. (a) Show that the cylindrical coordinate system is Orthogonal Curvilinear Coordinate System.
- (b) Express the vector  $\vec{f} = z \hat{i} - 2x \hat{j} + y \hat{k}$  in cylindrical coordinates and find  $f_\rho, f_\phi, f_z$



8. (a) Solve  $\frac{dx}{x(y^2 - z^2)} = \frac{dy}{y(z^2 - x^2)} = \frac{dz}{z(x^2 - y^2)}$

(b) Solve  $\frac{dx}{x^2 - yz} = \frac{dy}{y^2 - zx} = \frac{dz}{z^2 - xy}$

OR

9. (a) Express the vector  $\vec{f} = 3x\hat{i} - 2yz\hat{j} + x^2z\hat{k}$  in cylindrical coordinates and find  $f_\rho, f_\phi, f_z$

(b) Express the vector  $\vec{f} = x\hat{i} - y\hat{j} + z\hat{k}$  in spherical coordinates and find  $f_r, f_\theta, f_\phi$

PART - D

Answer **any two full** questions.

2x10=20

10. (a) Form the partial differential equation by eliminating the arbitrary functions  $z = f(x+ay) + g(x-ay)$

(b) Solve  $p(1+q) = zq$

OR

11. (a) Solve  $[D^2 - 2DD' + (D')^2]z = e^{x+2y}$

(b) Solve  $p+q = \sin x + \sin y$

12. (a) Find the complete integral of  $px+qy=pq$  by Charpit's method

(b) Solve  $[D^2 - 2DD' + (D')^2]z = 12xy$

OR

13. (a) A tightly stretched string with fixed end points  $x=0$  and  $x=1$  is initially in a position given by  $y = y_0 \sin^3\left(\frac{\pi x}{1}\right)$ . If it is released from rest from this position, find the displacement  $y(x, t)$ .

(b) Solve  $\frac{\partial u}{\partial t} = 16 \frac{\partial^2 u}{\partial x^2}$  subjected to the conditions.

(i)  $u(0, t) = 0, u(1, t) = 0$  for all  $t$

(ii)  $u(x, 0) = x^2 - x, 0 \leq x \leq 1$